

New localization method of $U(1)$ gauge vector field on flat branes in (asymptotic) AdS_5 spacetime

Zhen-Hua Zhao^{1, a}, Qun-Ying Xie,^{b, c} Yuan Zhong^{c, d}

^a*Department of Applied Physics, Shandong University of Science and Technology, Qingdao, 266590 People's Republic of China*

^b*School of Information Science and Engineering, Lanzhou University, Lanzhou 730000, People's Republic of China*

^c*Institute of Theoretical Physics, Lanzhou University, Lanzhou 730000, People's Republic of China*

^d*IFAE, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain*

E-mail: zhaozh09@lzu.edu.cn, xieqy@lzu.edu.cn, yzhong@ifae.es

ABSTRACT: It is well known that the $U(1)$ gauge vector field, with the standard five-dimensional (5D) action, cannot be localized on Randall-Sundrum-like braneworlds with an infinite extra dimension. In this paper, we propose a modified 5D action to localize $U(1)$ gauge vector field on flat branes with an infinite or finite extra dimension. The localization method is realized by adding a dynamical mass term into the standard 5D action of the vector field, which is proportional to the 5D scalar curvature. It is shown that the vector zero mode is localizable if the 5D spacetime is (asymptotic) AdS_5 . Moreover, the massive tachyonic modes can be excluded.

KEYWORDS: Extra Dimensions, Braneworld, Localization of Vector Field

¹Corresponding author

Contents

1	Introduction	1
2	Localization method of vector field on the branes	2
3	Excluding 4D vector tachyonic modes	5
4	Conclusion	6

1 Introduction

The braneworld theories have received a lot of attentions since the success of the Randall-Sundrum (RS) thin brane models [1, 2]. In braneworld theories, the localization of gravity and all kinds of matter fields on the brane is always an important issue. It is well known that the four-dimensional graviton can be localized in the RS thin brane scenario [1, 3] and in thick brane scenarios [4–10]. The localization of real scalar field is the same as gravity in general relativity [11], but may be different from or even opposite to gravity in some modified gravity [12, 13]. By introducing the usual Yukawa coupling between the background scalar field and the fermion field, the fermion can also be localized on the brane generated by an odd scalar field [11, 14–19]. However, if the brane is generated by an even scalar field, we need to introduce new localization mechanism in order to localize fermion on the brane [20].

In five-dimensional (5D) spacetime, the $U(1)$ gauge field $\mathcal{A}_M$ with the usual action

$$S \sim \int d^5x \sqrt{-g} \mathcal{F}_{MN} \mathcal{F}^{MN}, \quad (1.1)$$

where $\mathcal{F}_{MN} = \partial_M \mathcal{A}_N - \partial_N \mathcal{A}_M$ is the field strength, can be localized on the brane in some special braneworld models, for example, in the standing wave braneworld model [21], in the braneworld models with finite extra dimension [22–26], and in a 6D model [27]. But it cannot be localized in a RS-like braneworld model with an infinite extra dimension [11, 28, 29].

In order to localize $U(1)$ gauge field on branes in 5D RS-like models with an infinite extra dimension, the typical localization mechanism is to reform the action (1.1). In the thin brane scenario, many ideas were proposed for this issue [30–33]. In ref. [30], the author added a topological term and a 3-form gauge potential into the action (1.1). In ref. [31], a bulk mass term of the vector and a coupling between the vector potential and the brane were introduced. In ref. [32], the action (1.1) was changed to

$$S \sim \int d^5x \sqrt{-g} e^{2\alpha(y)} \mathcal{F}_{MN} \mathcal{F}^{MN}, \quad (1.2)$$

where $e^{2\alpha(y)}$ is the warp factor. In this model the vector zero mode can be localized on the negative tension brane. In ref. [33], gauge field kinetic terms induced by localized fermions were added into the gauge field action.

In order to localize $U(1)$ vector fields in a thick brane models, Kehagias and Tamvakis (KT) proposed a coupling between the gauge field and an extra dilaton field. The KT mechanism has been applied in many different brane-world scenarios [34–40]. Recently, Chumbes, Holf da Silva and Hott (CHH) proposed a new mechanism [29], in which gauge and tensor fields directly couple to a functional of the background scalar field. By introducing a Stueckelberg-like action, Vaquera-Araujo and Corradini realized the localization of vector in thick brane model[41].

In this paper, we propose a new localization method to localize $U(1)$ gauge field on the brane. We assume the 5D gauge field has a dynamic mass term, which is proportional to the 5D scalar curvature. Our localization method can be used both in thin and thick braneworld models with infinite or finite extra dimension. To localize gauge field zero mode, the only assumption we need is that the 5D spacetime is (asymptotic) AdS_5 . With the same assumption, we can further prove that there is no tachyonic mode in the KK spectrum.

The paper is structured as follows. In section 2, we show the setup of our localization method of gauge field on the brane and prove that the zero mode is localizable under the assumption of that the five-dimensional spacetime is (asymptotic) AdS_5 . Further, in section 3, we prove that tachyon modes can be excluded. We conclude our results in the last section.

2 Localization method of vector field on the branes

The line-element describing a flat (Minkowski) braneworld embedded in five-dimensional spacetime is assumed to be [1]

$$ds^2 = \mathcal{G}_{MN} dx^M dx^N = e^{2\alpha(y)} \eta_{\mu\nu} dx^\mu dx^\nu + dy^2, \quad (2.1)$$

where $e^{2\alpha(y)}$ is the warp factor, $\alpha(y)$ is a function of the extra dimension y , $\mathcal{G}_{MN}$ is the metric of 5D bulk spacetime and $M, N = 0, 1, 2, 3, 4$ stand for the bulk coordinate indices, and the Minkowski metric $\eta_{\mu\nu}$ on branes with signature $(-1, +1, +1, +1)$ and $\mu, \nu = 0, 1, 2, 3$ correspond to the brane coordinate indices.

In order to localize vector fields on branes, we introduce a dynamical mass term and the action reads

$$S = \int d^4x dy \sqrt{-\mathcal{G}} \left(-\frac{1}{4} \mathcal{G}^{MN} \mathcal{G}^{RS} \mathcal{F}_{MR} \mathcal{F}_{NS} - \frac{1}{2} \mathcal{M}^2 \mathcal{G}^{MN} \mathcal{A}_M \mathcal{A}_N \right), \quad (2.2)$$

where $\mathcal{M}^2$ is a function of the 5D scalar curvature. Here, we set

$$\mathcal{M}^2 = -\frac{1}{16} \mathcal{R}, \quad (2.3)$$

With the metric in Eq. (2.1), the scalar curvature is given by

$$\mathcal{R} = -4(5\alpha'^2 + 2\alpha''). \quad (2.4)$$

In this paper, the prime stands for the derivative with respect to y . Because the brane is flat so $\mathcal{R}$ is only a function of extra-dimension coordinate y .

Here, we parametrize $\mathcal{A}_M$ in the following way [31, 42, 43]:

$$\mathcal{A}_M = (\hat{\mathcal{A}}_\mu + \partial_\mu \phi, \mathcal{A}_4), \quad (2.5)$$

where $\hat{\mathcal{A}}_\mu$ is the transverse component which satisfies the transverse condition $\partial_\mu \hat{\mathcal{A}}^\mu = 0$, and ϕ is the longitudinal component. Substituting Eq. (2.5) into the action (2.2), with the transverse condition, one can find that the transverse vector $\hat{\mathcal{A}}_\mu$ decouples from the scalar fields ϕ and $\mathcal{A}_4$, and the action (2.2) can be split into two parts,

$$S = S_V(\hat{\mathcal{A}}_\mu) + S_S(\hat{\mathcal{A}}_4, \phi), \quad (2.6)$$

where [44, 45]

$$S_V = \int d^4x dy \sqrt{-\mathcal{G}} \left(-\frac{1}{4} \hat{\mathcal{F}}_{\lambda\mu} \hat{\mathcal{F}}_{\nu\rho} \mathcal{G}^{\lambda\nu} \mathcal{G}^{\mu\rho} - \frac{1}{2} \partial_4 \hat{\mathcal{A}}_\mu \partial_4 \hat{\mathcal{A}}_\nu \mathcal{G}^{44} \mathcal{G}^{\mu\nu} - \frac{1}{2} \mathcal{M}^2 \hat{\mathcal{A}}_\mu \hat{\mathcal{A}}_\nu \mathcal{G}^{\mu\nu} \right), \quad (2.7)$$

$$S_S = \int d^4x dy \sqrt{-\mathcal{G}} \left(-\frac{1}{2} \partial_4 (\partial_\mu \phi) \partial_4 (\partial_\nu \phi) \mathcal{G}^{44} \mathcal{G}^{\mu\nu} - \frac{1}{2} \mathcal{M}^2 \partial_\mu \phi \partial_\nu \phi \mathcal{G}^{\mu\nu} - \frac{1}{2} \partial_\mu \mathcal{A}_4 \partial_\nu \mathcal{A}_4 \mathcal{G}^{44} \mathcal{G}^{\mu\nu} - \frac{1}{2} \mathcal{A}_4 \mathcal{A}_4 \mathcal{M}^2 \mathcal{G}^{44} + \partial_\mu \mathcal{A}_4 \partial_4 (\partial_\nu \phi) \mathcal{G}^{44} \mathcal{G}^{\mu\nu} \right), \quad (2.8)$$

and $\hat{\mathcal{F}}_{\mu\nu} = \partial_\mu \hat{\mathcal{A}}_\nu - \partial_\nu \hat{\mathcal{A}}_\mu$.

Because we focus on the localization of vector field, and the scalar sector had been discussed in Ref. [43], therefore we will not discuss the localization of the scalar particles in this paper. The transverse vector field can be decomposed as

$$\hat{\mathcal{A}}_\mu(x, y) = \sum_n A_\mu^{(n)}(x) \chi_n(y), \quad (2.9)$$

where $A_\mu^{(n)}(x)$ is the 4D vector KK mode and $\chi_n(x)$ is called as the KK wave function [46], here and the following “ Σ_n ” is a shorthand for the summation and integration over discrete and continuum KK modes.

By means of the KK decomposition (2.9), the action (2.7) can be further reduced to

$$S_V = -\frac{1}{4} \sum_n \int dy \chi_n^2 \int d^4x (\eta^{\mu\nu} \eta^{\lambda\rho} F_{\mu\lambda}^{(n)} F_{\nu\rho}^{(n)} + 2m_n^2 \eta^{\mu\nu} A_\mu^{(n)} A_\nu^{(n)}), \quad (2.10)$$

where $F_{\mu\nu}^{(n)} = \partial_\mu A_\nu^{(n)} - \partial_\nu A_\mu^{(n)}$ is the four-dimensional vector field strength tensor. In order to obtain the action (2.10), the KK wave function $\chi_n(y)$ is required to satisfy the following equation

$$-\partial_y (e^{2\alpha} \partial_y \chi_n) + \chi_n e^{2\alpha} \mathcal{M}^2 = m_n^2 \chi_n, \quad (2.11)$$

and the orthonormalization condition

$$\int \chi_m(y) \chi_n(y) dy = 0, \quad (m \neq n). \quad (2.12)$$

From the actions (2.10), the localization of the 4D vector KK mode requires

$$I \equiv \int_{-\infty}^{+\infty} \chi_n^2(y) dy < \infty. \quad (2.13)$$

By using the following field transformation

$$\chi_n = e^{-\alpha} \tilde{\chi}_n, \quad (2.14)$$

Equation. (2.11) can be rewritten as

$$-\tilde{\chi}_n'' + (\alpha'' + \alpha'^2 + \mathcal{M}^2 - e^{-2\alpha} m_n^2) \tilde{\chi}_n = 0. \quad (2.15)$$

For the vector zero mode, $m_0 = 0$, Eq. (2.15) can be written as

$$-\tilde{\chi}_0'' + (\alpha'' + \alpha'^2 + \mathcal{M}^2) \tilde{\chi}_0 = 0. \quad (2.16)$$

Substituting Eqs. (2.3) and (2.4) into Eq. (2.16), we can obtain

$$-\tilde{\chi}_0'' + \left(\frac{3\alpha''}{2} + \frac{9\alpha'^2}{4}\right) \tilde{\chi}_0 = 0, \quad (2.17)$$

which can be further factorized as

$$\left(-\frac{d}{dy} - \frac{3}{2}\alpha'\right) \left(\frac{d}{dy} - \frac{3}{2}\alpha'\right) \tilde{\chi}_0 = 0. \quad (2.18)$$

The solution of the above equation is

$$\tilde{\chi}_0(y) = c_0 e^{\frac{3}{2}\alpha(y)}, \quad (2.19)$$

and further one can get

$$\chi_0 = e^{-\alpha} \tilde{\chi}_0 = c_0 e^{\frac{1}{2}\alpha}. \quad (2.20)$$

With the above solution, the integration (2.13) reads

$$I = \int_{-\infty}^{+\infty} \chi_0^2 dy = c_0^2 \int_{-\infty}^{+\infty} e^{\alpha} dy. \quad (2.21)$$

Because the integrand in Eq. (2.21) is continuous at the interval $y \in (-\infty, +\infty)$, the convergence of the above integrations is determined by the asymptotic behaviors of integrands at the infinity.

Now, we prove that if the five-dimensional spacetime is asymptotic AdS_5 , the vector zero mode is always localizable. The condition of asymptotic AdS_5 means that, when $y \rightarrow \pm\infty$, the warp factor has the following asymptotic behavior

$$\alpha(y)|_{y \rightarrow \pm\infty} \rightarrow -k|y|, \quad (2.22)$$

where k is a positive constant and it usually relates with the five-dimensional fundamental scale in RS-like braneworld scenarios. With this condition, the asymptotic behavior of integrand in Eq. (2.21) reads as

$$e^{\alpha}|_{y \rightarrow \pm\infty} \rightarrow e^{-k|y|}. \quad (2.23)$$

Therefore, the integration (2.21) is convergent, namely, the 4D vector zero mode is localizable. If the extra dimension is finite, the conclusion is also valid.

Our conclusion can be easily verified in the RS-II model [3], in which $\alpha(y) = -k|y|$. Substituting it into Eq. (2.21), we obtain

$$I = 2c_0^2 \int_0^{+\infty} e^{-ky} dy = 2c_0^2/k. \quad (2.24)$$

Normalizing I results in $c_0 = \sqrt{k/2}$. Therefore, the vector zero mode is localizable in RS-II model.

3 Excluding 4D vector tachyonic modes

Further, we will discuss if this new localization method can exclude the tachyonic modes of 4D vector field. To discuss the massive modes and mass spectrum, it is more convenient to work with the conformal metric:

$$ds^2 = e^{2\alpha(z)} (\eta_{\mu\nu} dx^\mu dx^\nu + dz^2). \quad (3.1)$$

The form of dynamic mass with above metric (3.1) is

$$\mathcal{M}^2 = -\frac{1}{16}\mathcal{R} = \frac{1}{4}e^{-2\alpha}(3\dot{\alpha}^2 + 2\ddot{\alpha}). \quad (3.2)$$

where the symbol “ $\cdot$ ” means the derivative with respect to z .

Making use of the KK decomposition

$$\hat{\mathcal{A}}_\mu(x, z) = \sum_n A_\mu^{(n)}(x) \chi_n(z), \quad (3.3)$$

and the transformation

$$\chi_n(z) = \rho_n(z) e^{-\alpha/2}, \quad (3.4)$$

the action (2.7) reduces to

$$S_V = -\frac{1}{4} \sum_n \int dz \rho_n(z)^2 \int d^4x (F_{\mu\nu}^{(n)} F^{(n),\mu\nu} + 2m_n^2 A_\mu^{(n)} A^{(n),\mu}). \quad (3.5)$$

The localization of KK modes means

$$I \equiv \int_{-\infty}^{+\infty} \rho_n^2(y) dy < \infty. \quad (3.6)$$

At the same time, the function $\rho(z)$ is required to satisfy a Schrödinger-like equation

$$(-\partial_z^2 + V(z)) \rho_n(z) = m_n^2 \rho_n(z) \quad (3.7)$$

and the orthonormalization condition

$$\int \rho_m(z) \rho_n(z) dz = 0 \quad (m \neq n). \quad (3.8)$$

The effective potential is given by,

$$V(z) = \frac{1}{2}\ddot{\alpha}(z) + \frac{1}{4}\dot{\alpha}(z)^2 + e^{2\alpha}\mathcal{M}^2. \quad (3.9)$$

Substituting the expression of $\mathcal{M}^2$ in (3.2) into Eq. (3.7), we obtain

$$V(z) = \ddot{\alpha}(z) + \dot{\alpha}(z)^2. \quad (3.10)$$

With above potential Eq. (3.7) can be further factored as

$$(-\partial_z - \dot{\alpha})(\partial_z - \dot{\alpha})\rho_n(z) = m_n^2\rho_n(z). \quad (3.11)$$

This is the supersymmetry quantum mechanics form of the Schrödinger equation (3.7) [47], which guarantees the positivity of m_n^2 . Therefore, the tachyonic vector modes are excluded. This result is independent of the asymptotic behavior of spacetime.

4 Conclusion

In this paper, we proposed a new localization method of $U(1)$ gauge field. In our method a dynamical mass term was added into the 5D action of the vector field. The dynamical mass term is proportional to the 5D scalar curvature. It was shown that, if the brane is embedded in a 5D AdS_5 spacetime, then the vector zero mode is localized on the brane. Moreover, we also proved that there is no vector tachyonic mode with our method.

Acknowledgments

We thank Prof. Y.-X. Liu for helpful discussion. Z.-H. Zhao is supported by the National Natural Science Foundation of China (Grant No. 11305095), the Natural Science Foundation of Shandong Province, China (Grant No. ZR2013AQ016), and Scientific Research Foundation of Shandong University of Science and Technology for Recruited Talents (Grant No. 2013RCJJ026). Q.Y. Xie is supported by the National Natural Science Foundation of China (Grant No. 11375075). Y. Zhong is supported by the scholarship granted by the Chinese Scholarship Council (CSC).

References

- [1] L. Randall and R. Sundrum, *A large mass hierarchy from a small extra dimension*, Phys. Rev. Lett. **83** (1999) 3370, [[hep-ph/9905221](#)].
- [2] J. Garriga and T. Tanaka, *Gravity in the randall-sundrum brane world*, Phys. Rev. Lett. **84** (2000) 2778, [[hep-th/9911055](#)].
- [3] L. Randall and R. Sundrum, *An alternative to compactification*, Phys. Rev. Lett. **83** (1999) 4690, [[hep-th/9906064](#)].
- [4] M. Gremm, *Four-dimensional gravity on a thick domain wall*, Phys. Lett. B **478** (2000) 434, [[hep-th/9912060](#)].
- [5] D. Bazeia and A. R. Gomes, *Bloch brane*, JHEP **05** (2004) 012, [[hep-th/0403141](#)].

- [6] N. Barbosa-Cendejas and A. Herrera-Aguilar, *4d gravity localized in non z_2 -symmetric thick branes*, JHEP **10** (2005) 101, [[hep-th/0511050](#)].
- [7] N. Barbosa-Cendejas and A. Herrera-Aguilar, *Localization of 4d gravity on pure geometrical thick branes*, Phys. Rev. D **73** (2006) 084022, [[hep-th/0603184](#)].
- [8] K. Farakos, G. Koutsoumbas, and P. Pasipoularides, *Graviton localization and newton's law for brane models with a non-minimally coupled bulk scalar field*, Phys. Rev. D **76** (2007) 064025, [[arXiv:0705.2364](#)].
- [9] G. German, A. Herrera-Aguilar, D. Malagon-Morejon, R. R. Mora-Luna, and R. da Rocha, *A de sitter tachyon thick braneworld*, JCAP **1302** (2013) 035, [[arXiv:1210.0721](#)].
- [10] A. Herrera-Aguilar, D. Malagon-Morejon, R. R. Mora-Luna, and I. Quiros, *Thick braneworlds generated by a non-minimally coupled scalar field and a gauss-bonnet term: conditions for localization of gravity*, Class. Quant. Grav. **29** (2012) 035012, [[arXiv:1105.5479](#)].
- [11] B. Bajc and G. Gabadadze, *Localization of matter and cosmological constant on a brane in anti de sitter space*, Phys. Lett. B **474** (2000) 282, [[hep-th/9912232](#)].
- [12] Y. Zhong, Y.-X. Liu, and K. Yang, *Tensor perturbations of $f(r)$ -branes*, Phys. Lett. B **699** (2011) 398, [[arXiv:1010.3478](#)].
- [13] K. Yang, Y.-X. Liu, Y. Zhong, X.-L. Du, and S.-W. Wei, *Gravity localization and mass hierarchy in scalar-tensor branes*, Phys. Rev. D **86** (2012) 127502, [[arXiv:1212.2735](#)].
- [14] V. A. Rubakov and M. E. Shaposhnikov, *Do we live inside a domain wall?*, Phys. Lett. B **125** (1983) 136.
- [15] S. Randjbar-Daemi and M. E. Shaposhnikov, *Fermion zero-modes on brane-worlds*, Phys. Lett. B **492** (2000) 361, [[hep-th/0008079](#)].
- [16] C. Ringeval, P. Peter, and J.-P. Uzan, *Localization of massive fermions on the brane*, Phys. Rev. D **65** (2002) 044016, [[hep-th/0109194](#)].
- [17] R. Koley and S. Kar, *Scalar kinks and fermion localization in warped spacetimes*, Class. Quant. Grav. **22** (2005) 753, [[hep-th/0407158](#)].
- [18] A. Melfo, N. Pantoja, and J. D. Tempo, *Fermion localization on thick branes*, Phys. Rev. D **73** (2006) 044033, [[hep-th/0601161](#)].
- [19] Y.-X. Liu, L.-D. Zhang, L.-J. Zhang, and Y.-S. Duan, *Fermions on thick branes in background of sine-gordon kinks*, Phys. Rev. D **78** (2008) 065025, [[arXiv:0804.4553](#)].
- [20] Y.-X. Liu, Z.-G. Xu, F.-W. Chen, and S.-W. Wei, *New localization mechanism of fermions on braneworlds*, Phys. Rev. D **89** (2014) 086001, [[arXiv:1312.4145](#)].
- [21] M. Gogberashvili, P. Midodashvili, and L. Midodashvili, *Localization of gauge bosons in the 5d standing wave braneworld*, Phys. Lett. B **707** (2012) 169, [[arXiv:1105.1866](#)].
- [22] Y.-X. Liu, L.-D. Zhang, S.-W. Wei, and Y.-S. Duan, *Localization and mass spectrum of matters on weyl thick branes*, JHEP **08** (2008) 041, [[arXiv:0803.0098](#)].
- [23] Y.-X. Liu, Z.-H. Zhao, S.-W. Wei, and Y.-S. Duan, *Bulk matters on symmetric and asymmetric de sitter thick branes*, J. Cosmol. Astropart. Phys. **0902** (2009) 003, [[arXiv:0901.0782](#)].
- [24] Y.-X. Liu, C.-E. Fu, H. Guo, and H.-T. Li, *Deformed brane with finite extra dimension*, Phys. Rev. D **85** (2012) 084023, [[arXiv:1102.4500](#)].

- [25] H. Guo, A. Herrera-Aguilar, Y.-X. Liu, D. Malagon-Morejon, and R. R. Mora-Luna, *Localization of bulk matter fields on a pure de sitter thick braneworld*, Phys. Rev. D **87** (2013) 095011, [[arXiv:1103.2430](#)].
- [26] A. Herrera-Aguilar, A. D. Rojas, and E. Santos-Rodriguez, *Localization of gauge fields in a tachyonic de sitter thick braneworld*, Eur. Phys. J. C **74** (2014) 2770, [[arXiv:1401.0999](#)].
- [27] I. Oda, *Localization of matters on a string - like defect*, Phys. Lett. B **496** (2000) 113, [[hep-th/0006203](#)].
- [28] A. Pomarol, *Gauge bosons in a five-dimensional theory with localized gravity*, Phys. Lett. B **486** (2000) 153, [[hep-ph/9911294](#)].
- [29] A. Chumbes, J. Hoff da Silva, and M. Hott, *A model to localize gauge and tensor fields on thick branes*, Phys. Rev. D **85** (2012) 085003, [[arXiv:1108.3821](#)].
- [30] I. Oda, “A new mechanism for trapping of photon.” 2001.
- [31] K. Ghoroku and A. Nakamura, *Massive vector trapping as a gauge boson on a brane*, Phys. Rev. D **65** (2002) 084017, [[hep-th/0106145](#)].
- [32] M. Giovannini, *Vector field localization and negative tension branes*, Phys. Rev. D **65** (2002) 124019, [[hep-th/0204235](#)].
- [33] R. Guerrero, A. Melfo, N. Pantoja, and R. O. Rodriguez, *Gauge field localization on brane worlds*, Phys. Rev. D **81** (2010) 086004, [[arXiv:0912.0463](#)].
- [34] W. Cruz, M. Tahim, and C. Almeida, *Gauge field localization on a dilatonic deformed brane*, Phys. Lett. B **686** (2010) 259.
- [35] G. Alencar, R. Landim, M. Tahim, C. Muniz, and R. Costa Filho, *Antisymmetric tensor fields in randall sundrum thick branes*, Phys. Lett. B **693** (2010) 503, [[arXiv:1008.0678](#)].
- [36] W. Cruz, A. R. Lima, and C. Almeida, *Gauge field localization on the bloch brane*, Phys. Rev. D **87** (2013) 045018, [[arXiv:1211.7355](#)].
- [37] C.-E. Fu, Y.-X. Liu, and H. Guo, *Bulk matter fields on two-field thick branes*, Phys. Rev. D **84** (2011) 044036, [[arXiv:1101.0336](#)].
- [38] W. Cruz, M. Tahim, and C. Almeida, *Results in kalb-ramond field localization and resonances on deformed branes*, Europhys.Lett. **88** (2009) 41001, [[arXiv:0912.1029](#)].
- [39] H. Christiansen, M. Cunha, and M. Tahim, *Exact solutions for a maxwell-kalb-ramond action with dilaton: Localization of massless and massive modes in a sine-gordon brane-world*, Phys. Rev. D **82** (2010) 085023, [[arXiv:1006.1366](#)].
- [40] W. Cruz, R. Maluf, and C. Almeida, *Kalb-ramond field localization on the bloch brane*, [arXiv:1303.1096](#).
- [41] C. A. Vaquera-Araujo and O. Corradini, *Localization of abelian gauge fields on thick branes*, [arXiv:1406.2892](#).
- [42] B. Batell and T. Gherghetta, *Localized $u(1)$ gauge fields, millicharged particles, and holography*, Phys. Rev. D **73** (2006) 045016.
- [43] G. Alencar, R. Landim, M. Tahim, and R. N. C. Filho, *Gauge field localization on the brane through geometrical coupling*, [arXiv:1409.4396](#).
- [44] K. Peeters, *A field-theory motivated approach to symbolic computer algebra*, Comput.Phys.Commun. **176** (2007) 550, [[cs/0608005](#)].

- [45] K. Peeters, *Introducing cadabra: A symbolic computer algebra system for field theory problems*, [hep-th/0701238](#).
- [46] E. Ponton, *Tasi 2011: Four lectures on tev scale extra dimensions*, [arXiv:1207.3827](#).
- [47] A. A. Andrianov, V. A. Andrianov, P. Giacconi, and R. Soldati, *Domain wall generation by fermion self-interaction and light particles*, JHEP **07** (2003) 063, [[hep-ph/0305271](#)].